

Leafminer Parasitoid *Opius dissitus* Muesebeck (Insecta: Hymenoptera: Braconidae)¹

Jian Li, Dakshina Seal, and Gary Leibe²

Introduction

The wasp *Opius dissitus* Muesebeck is a solitary, larva-pupal, Hymenopteran endoparasitoid of *Liriomyza* leafminers. Several studies report that *O. dissitus* were reared from *Liriomyza* (Diptera: Agromyzidae) leafminers infesting plant leaves of celery, tomato, potato, beans, etc. (unpublished data, Li et al.; Stegmaier 1972).

Distribution

Opius dissitus is reported from Africa, Asia, Europe, and North America (Bordat et al. 1995a; Petcharat et al. 2002; Stegmaier 1972; Neuenschwander 1987).

Description and Life Cycle

Adults

The adult *O. dissitus* are black in color. The antennae are black and thin, and almost the same length as their body.

Eggs

Opius dissitus females lay their eggs directly inside the late stage *Liriomyza* larvae bodies. The average size of the egg is about 0.28 mm (Bordat et al. 1995a).



Figure 1. Dorsal view of an adult *Opius dissitus* Muesebeck, an endoparasitoid of *Liriomyza* leafminers.

Credits: Jian Li, UF/IFAS

Larvae

The *O. dissitus* larvae develop inside the *Liriomyza* leafminer larvae. However, the parasitized leafminer larvae still consume the tissue of plant leaves until their pupation. The *O. dissitus* larvae develop through two instars inside the leafminer larvae (Bordat et al. 1995a), and eventually kill the leafminer in the pupal stage. The mature *O. dissitus* larvae then pupate inside the leafminer pupae. The optimal

1. This document is EENY 501, one of a series of the Department of Entomology and Nematology, UF/IFAS Extension. Original publication date October 2011. Reviewed August 2018. Visit the EDIS website at <http://edis.ifas.ufl.edu>. This document is also available on the Featured Creatures website at <http://entnemdept.ifas.ufl.edu/creatures/>.

2. Jian Li; Dakshina Seal; and Gary Leibe; Department of Entomology and Nematology, UF/IFAS Tropical Research and Education Center, Lake Alfred, FL 33850.



Figure 2. Early stage pupa of *Opius dissitus* Muesebeck, an endoparasite of *Liriomyza* leafminers.
Credits: Jian Li, UF/IFAS



Figure 3. Early stage pupa of *Opius dissitus* Muesebeck, an endoparasite of *Liriomyza* leafminers.
Credits: Jian Li, UF/IFAS

temperature for *O. dissitus* development on host of *L. trifolii* is reported as 25°C–30°C (Bordat et al. 1995b).

Pupae

Opius dissitus early stage pupae are yellow and have red eyes, while the mature stage pupae are black in color. *Opius dissitus* adults emerge out of the leafminer pupae. One adult emerges from a single parasitized pupa (unpublished data Li et al.)

Hosts

In Florida, *Opius dissitus* larvae were found and collected, then reared to adults, from the leaves of several crops

infested by *Liriomyza* leafminers (Stegmaier 1972; Schuster and Wharton 1993). Important *Liriomyza* economic pest species include the pea leafminer, *L. huidobrensis* (Blanchard); the vegetable leafminer, *L. sativae* Blanchard; and the American serpentine leafminer, *L. trifolii* (Burgess).

Economic Importance

Opius dissitus is a potential biological control agent for *Liriomyza* leafminers on vegetable and ornamental plants. *Opius dissitus* was found to be the most abundant parasitoid (63% of all the parasitoids) of *L. trifolii* on snap bean crops in south Florida, and the seasonal density of *O. dissitus* had a similar pattern with *L. trifolii* (unpublished data Li et al.). *Opius dissitus* was also found to be one of the major hymenopteran parasitoids of *Liriomyza* leafminer on tomato crops (Schuster and Wharton 1993). Pettitt (2004) reported that *O. dissitus* was reared and released to control *Liriomyza* leafminers at the Walt Disney World Resort.

Selected References

- Bordat D. 1995a. "Morphometric, biological and behavioral differences between (Hym., Eulophidae) and (Hym., Braconidae) parasitoids of (Dipt., Agromyzidae)." *Journal of Applied Entomology* 119: 423–427.
- Bordat D. 1995b. "Influence of temperature on (Hym., Braconidae), a parasitoid of (Dipt: Agromyzidae)." *Entomophaga* 40: 119–124.
- Neuenschwander P. 1987. "Introduction of exotic parasitic wasps for the control of (Diptera: Agromyzidae) in Senegal." *Tropical Pest Management* 33: 290–297.
- Petcharat J. 2002. "Larval parasitoids of agromyzid leafminer genus in the southern Thailand: species and their host plants." *Songklanakarin Journal of Scientific Technology* 24: 467–472.
- Pettitt LF. 2004. "Rearing and release of (Hymenoptera: Braconidae) for biological control of leafminers at the Walt Disney World Resort. Symposium: Advances In Management For Agromyzid Leaf Miners." *Entomological Society of America Annual Meeting and Exhibition*.
- Schuster DJ, Wharton RA. 1993. "Hymenopterous parasitoids of leaf-mining spp. (Diptera: Agromyzidae) on tomato in Florida." *Environmental Entomology* 22: 1188–1191.
- Stegmaier CE. 1972. "Parasitic Hymenoptera bred from the family Agromyzidae (Diptera) with special reference to south Florida." *Florida Entomologist* 55: 273–282.