

Integrated Pest Management in Protected Structures I: Basic Principles and Scouting¹

Hugh A. Smith, Gary E. Vallad, and Bielinski M. Santos²

The fundamentals of managing pests in protected structures are very similar in many respects to managing pests in field crops. The grower must still develop a management plan, assembling pertinent information for identifying, monitoring and controlling specific pests before planting. Integration of several pest management strategies into a comprehensive program is still critical. Recognition of seasonal patterns and anticipation of associated pests are important for timing control measures. Unlike field pest management, conditions within a protected structure can be modified to a certain degree to prevent, delay, or even mitigate pest issues. However, it's important to recognize that conditions that discourage one group of pests can often favor another. Protected structures can be screened to physically exclude many insect pests (Figure 1), but screens also shelter insect pests inside the structure from enemies. Structures with non-porous tops like greenhouses (Figure 2) and high tunnels (Figure 3) can mitigate the effect of rain on many bacterial and fungal pathogens that require free moisture for infection and movement, but without proper ventilation these same structures will increase humidity to a degree that offsets these gains or encourages other diseases such as *Botrytis* and powdery mildew. While the use of soilless media can eliminate many detrimental soil-borne pests and pathogens that plague field production, some soil-borne pathogens produce spores that can still be problematic in a protected structure even when the pathogen source is outside the structure.



Figure 1. Screen house made with insect-exclusion material.
Credits: Bielinski Santos



Figure 2. Greenhouse.
Credits: Bielinski Santos

1. This document is ENY868, one of a series of the Entomology and Nematology Department, UF/IFAS Extension. Original publication date June 2013. Reviewed September 2019. Visit the EDIS website at <https://edis.ifas.ufl.edu> for the currently supported version of this publication.
2. Hugh A. Smith, assistant professor, Entomology and Nematology Department; Gary E. Vallad, associate professor, Department of Plant Pathology; and Bielinski M. Santos, former associate professor of horticulture, Vegetable and Fruit Crop Horticultural Program; UF/IFAS Gulf Coast Research and Education Center; UF/IFAS Extension, Gainesville, FL 32611.

The Institute of Food and Agricultural Sciences (IFAS) is an Equal Opportunity Institution authorized to provide research, educational information and other services only to individuals and institutions that function with non-discrimination with respect to race, creed, color, religion, age, disability, sex, sexual orientation, marital status, national origin, political opinions or affiliations. For more information on obtaining other UF/IFAS Extension publications, contact your county's UF/IFAS Extension office. U.S. Department of Agriculture, UF/IFAS Extension Service, University of Florida, IFAS, Florida A & M University Cooperative Extension Program, and Boards of County Commissioners Cooperating. Nick T. Place, dean for UF/IFAS Extension.



Figure 3. High tunnel with sides raised and ends open.

Credits: Hugh Smith

There are also greater limitations on the use of pesticides in protected structures. Depending on the type of structure, there may be fewer registered pesticides available compared to the array of options available for field production. Some materials cannot be applied in structures because of worker safety concerns. Other materials are not permitted because they leave pesticide residues. Depending on the crop and production strategy, production in a protected structure can take considerably longer than field production (Figure 4). These longer production periods sometimes present a problem because many pesticide labels have restricted application frequencies and maximum rates that were developed for commercial field production. The restrictions on pesticides that can be used and the potential for prolonged production mean that producers must plan carefully to ensure proper pesticide rotations so that pests in protected structures don't become resistant to pesticides. Fortunately, protected structures offer unique opportunities to use parasitic and predatory arthropods (Figure 5A–D) as well as other biological control strategies to offset the use of conventional pesticides.



Figure 4. Indeterminate tomatoes grown in a screen house. Indeterminate crops may have production seasons of several months. Rates and frequencies of applications for some insecticides are developed for the shorter field cycle, with the result that producers of indeterminate crops may run out of spray options.

Credits: Hugh Smith




Figure 5. A: The predatory mite *Amblyseius swirskii* feeding on whitefly nymphs, B: *Eretmocerus eremicus*, parasitic wasps that attack whitefly nymphs, C: the predatory mite *Phytoseiulus persimilis*, which aids in the control of spider mites, D: the predatory minute pirate bug *Orius insidiosus* feeding on thrips.

Credits: Lyle Buss

I. Prevention

As Henry de Bracton wrote, “An ounce of prevention is worth a pound of cure.” The key to successful pest management is to prevent the problem from occurring.

- Avoid production near obvious sources of infestation, such as fields in which the same crop or a related crop is being produced. Be sure to destroy crops from the previous season well in advance of production, and ensure that all residues are buried to destroy possible sources of bacterial, fungal, and viral pathogens. This also includes weeds and volunteer plants near the structure that can be sources of insect pests and plant pathogens.
- Sanitation is important and begins before planting. Many products including—alcohol, sodium hypochlorite, or ammonium-based materials—can be used to sanitize structures, pots, equipment, and other reused materials.
- Make plans to sterilize your growing medium. Treat soils for common soil-borne pests and pathogens, either by fumigation or other cultural methods (solarization, steam pasteurization, or flooding). If you re-use a soilless media, it's safest to sterilize it before you plant your new crop.
- It is equally important to start with plant materials (seed, transplants, and cuttings) that are healthy and free of pests and disease (Figure 6). “Seedborne” plant pathogens, as their name implies, are carried on seed, and there are many of them, so it is wise to purchase seeds only from reputable vendors. Seed can be treated with hot water or sodium hypochlorite to destroy pathogens

A large, rectangular tray filled with numerous small, green seedlings growing in a grid pattern. The seedlings are densely packed and appear to be in the early stages of growth. The tray is situated in a greenhouse or nursery setting, with other similar trays visible in the background. The lighting is bright, and the overall scene suggests a controlled environment for plant propagation.

- Many types of insect-exclusion material will prevent the entry of insects such as whiteflies, thrips, aphids, and mites into protected structures. Materials that reduce UV light have been successfully used to reduce damage by whiteflies, aphids, and thrips in protected structures in Israel and California. These insects apparently use UV light to navigate and are deterred from colonizing areas with insufficient UV light. The advantages of insect-excluding and UV-absorbing materials are reduced when walls and ceilings must be raised or opened for ventilation.

The success of any pest management strategy rests on timely identification of the pest or disease problem. Depending on the size of the operation and types of plants grown, the goal is to examine the majority of plants twice a week. If the operation is too large to examine every plant, follow a scouting plan that will ensure that different sections of production are examined each time. Basic scouting tools include a 10-40x hand lens, pocket knife, or pruning shears, and a scouting sheet (Figure 7). Scouting sheets are used to record the number of plants examined, pest or beneficial arthropods observed, and possible disease

[illegible]

A photograph of a cardboard box filled with various rolls of safety tape. The tapes are in different colors and patterns: orange, pink, yellow, black and yellow striped, blue, and white and blue striped. A long strip of orange tape is unrolled in front of the box, showing its length and the roll's core.

Integrated Pest Management in Protected Structures I: Basic Principles and Scouting 3



Figure 9. A tomato plant wilting because of *Fusarium* crown rot.
Credits: Gary Vallad



Figure 10. Squash leaves distorted from infection with Cucurbit leaf crumple virus, which is vectored by silverleaf whitefly.
Credits: Gary Vallad




Figure 11. Various plant diseases cause spotting or lesions. A: Late blight on tomato, B: downy mildew on cucumber, C: *Stemphylium* on tomato, D: pepper leaf infected with *Cercospora*.
Credits: Gary Vallad

Methods of scouting will vary according to the crop and the particular pest or disease, but here are some general considerations.

For diseases:

- Get familiar with plant structures and what constitutes a departure from normal growth.
- Know what environmental conditions and seasonal patterns favor which diseases, so you can better focus your scouting.
- Spores of fungal pathogens develop at night. They are more numerous and therefore easier to view within lesions first thing in the morning when it is light enough to see but before they have dispersed. Use your hand lens to spot the tiny spores.
- Be sure to check the underside of leaves because the initial symptoms of some diseases are more evident on the underside of leaves than the top.
- Most symptoms of viral pathogens are evident in the new growth of plants.
- Nicked or cut stems (from pruning) are ideal sites for *Botrytis* and other opportunistic pathogens to establish.
- Powdery mildews typically start in older foliage of plants, often when the plants begin to set fruit or approach

vegetative maturity. In many cases, the disease starts on the underside of leaves (Figure 12).



Figure 12. Powdery mildew on squash leaf.
Credits: Gary Vallad

- Symptoms of most vascular wilt diseases will be uneven on the plant (often one sided) (Figure 13) and on individual leaves. Look for a distinct discoloration of vascular tissues (Figure 14); it is quite unlike wilt caused by inadequate water.
- Sudden green wilts are often caused by a severe crown infection, by girdling, or by inadequate water.



Figure 13. One-sided foliar damage typical of vascular wilt diseases.
Credits: Gary Vallad



Figure 14. Discoloration of vascular tissue in tomato from *Fusarium* crown rot.
Credits: Gary Vallad

For insect or arthropod pests:

- Aphids can be found anywhere on a plant, but they prefer growing points (Figure 15A and B).




Figure 15. Aphids are serious pests of horticultural and ornamental crops grown in protected structures. A: Winged green peach aphid, B: infestation of melon aphids, including wingless adults and small nymphs. Note brownish parasitized aphid "mummy," C: aphid parasitoid.

Credits: Hugh Smith

- Whitefly adults settle on the underside of leaves, where they lay eggs (Figure 16A). Whitefly nymphs are only mobile immediately after leaving the egg; they move away from the hatched egg, settle down to feed, and do not move during subsequent nymphal development. For this

reason older whitefly nymphs are found on older leaves (Figure 16B).



Figure 16. A: Silverleaf whitefly adults, B: silverleaf whitefly nymphs. Note empty cast skins ("exuviae") from which adult whiteflies have emerged at lower right of picture.
Credits: James Castner and Lyle Buss

- Western flower thrips (Figure 17A) and other flower thrips are often hidden within flowers, where they feed on pollen. Blow or tap on the flower to dislodge the thrips and determine if they are present. Thrips puncture the plant cells and produce abrasions on the surface of leaves and flower petals. Thrips also deposit dark pellets of frass (excrement) in areas where they have fed, and this can be used to identify thrips damage even when thrips are not visible (Figure 17B).



Figure 17. A: Western flower thrips adult, B: feeding damage and frass left by Western flower thrips.
Credits: Lyle Buss and Hugh Smith

- Spider mites often establish on the underside of leaves (Figure 18A). They feed by puncturing plant cells, causing damage similar to thrips damage. Spider mites produce characteristic webbing (Figure 18 \B).



Figure 18. A: Twospotted spider mite, B: incipient feeding damage and webbing from twospotted spider mite infestation along mid-vein of eggplant leaf.
Credits: Lyle Buss and Yankai Li.

- Broad mites (Figure 19A) and cyclamen mites are much smaller than spider mites and are not easily detected with the naked eye. Broad mite feeding on pepper produces a

characteristic blistering or strappingness of leaves (Figure 19B). Examine upper leaves with a hand lens to spot broad mites.



Figure 19. A: Broad mite female and eggs on pepper leaf, B: distortion of pepper leaves due to broad mite feeding.
Credits: Hugh Smith

Yellow sticky cards are the most useful and commonly used monitoring tools to support a scouting program. Whiteflies, aphids, thrips, leafminers, leafhoppers, fungus gnats (Figure 20A), and shoreflies (Figure 20B) are all attracted to yellow sticky cards. Yellow sticky cards will trap primarily the adult stage of these insects because only the adults can fly. In the case of aphids, only alate (winged) adults will be trapped (Figure 15A). Some commercial biocontrol agents, such as parasitic wasps used to manage whiteflies, aphids, and other pests, are also attracted to yellow sticky cards (Figure 21). (Finding a limited number of parasitic wasps on yellow sticky cards in a greenhouse where they have been released indicates that the parasitoids are emerging and searching for hosts in significant numbers and should not be cause for concern.)



Figure 20. A: Fungus gnats and B: shore flies can become abundant in greenhouses and other protected structures when soil humidity is high.
Credits: Hugh Smith



Figure 21. Whitefly parasitoid on yellow sticky card.

Credits: Hugh Smith

Some growers use **sentinel plants** to reveal the presence of pests in protected structures. Eggplant has been used as a sentinel plant for whiteflies and green beans for spider mites. Sentinel plants may be effective when the pest is easier to detect on the sentinel plant than on the main crop. If you use sentinel plants, take care that they do not serve as a source of infestation for the main crop. Keep accurate scouting records in order to become familiar with pest cycles in a given structure and to evaluate the effectiveness of your control measures.