


Lyme Disease¹

P. E. Kaufman, P. G. Koehler, J. F. Butler, and A. Chaskopoulou²



This fact sheet is excerpted from *SP486: Pests in and around the Southern Home*, which is available from the UF/IFAS Extension Bookstore. <http://ifasbooks.ifas.ufl.edu/p-1222-pests-in-and-around-the-southern-home.aspx>

Lyme disease was first documented in the United States in 1975. The organism that causes Lyme disease is transmitted by the bite of a tick. Lyme disease is named for the town in Connecticut where some of the first cases occurred in the United States. Lyme disease cases have since been documented in about 43 states, with over 15,000 cases reported

annually between 1998 and 2004. Over 153,000 total cases were reported in the United States between 1993 and 2002. During 2002, 23,763 Lyme disease cases were reported, making this the peak year. During 2003-2005, 64,382 cases were reported. During the year of 2007, 27,444 cases of Lyme disease were reported, yielding a national average of 9.1 cases per 100,000 persons. Several cases of Lyme disease have been reported in Florida as well, corresponding to approximately 40 cases per year.

The primary vector of Lyme disease in the United States is the black-legged deer tick (Figures 1 and 2). Other closely related ticks are also known to transmit the disease. Lyme disease is maintained in wild rodent populations, on which the larval and nymphal ticks develop. These immature ticks pick up the disease organism when they suck the blood of infected rodents. The nymphal and adult ticks then seek a larger host, such as deer or humans, to obtain their final blood meal and transmit the disease when they feed.

Symptoms

The first sign of Lyme disease in 70%-80% of patients is a red circular rash, called an erythema migrans, around the puncture mark made where the tick pierced the skin. This rash appears after a three-to-30 day delay. The most common shape of the rash is an oval 2- to 3-inches in diameter that usually lasts about four weeks. The center of the rash occasionally will lighten resulting in a bull's-eye appearance.

1. This document is ENY-209, one of a series of the Department of Entomology and Nematology, UF/IFAS Extension. Original publication date October 1998. Revised October 2013. Reviewed September 2017. Visit the EDIS website at <http://edis.ifas.ufl.edu>.
2. P. E. Kaufman, professor; P. G. Koehler, professor; J. F. Butler, professor (retired); and A. Chaskopoulou, former graduate student; Department of Entomology and Nematology, UF/IFAS Extension, Gainesville, FL 32611.

The use of trade names in this publication is solely for the purpose of providing specific information. UF/IFAS does not guarantee or warranty the products named, and references to them in this publication do not signify our approval to the exclusion of other products of suitable composition.

The Institute of Food and Agricultural Sciences (IFAS) is an Equal Opportunity Institution authorized to provide research, educational information and other services only to individuals and institutions that function with non-discrimination with respect to race, creed, color, religion, age, disability, sex, sexual orientation, marital status, national origin, political opinions or affiliations. For more information on obtaining other UF/IFAS Extension publications, contact your county's UF/IFAS Extension office.

U.S. Department of Agriculture, UF/IFAS Extension Service, University of Florida, IFAS, Florida A & M University Cooperative Extension Program, and Boards of County Commissioners Cooperating. Nick T. Place, dean for UF/IFAS Extension.

The rash does not itch but may feel warm to the touch. Some patients might develop additional lesions in other parts of their body. Flu-like symptoms may also develop that often include aches, fever, fatigue, muscle pain, joint pain, and headache. Arthritis, cardiac disease, and neurologic disorders may develop if the disease is not properly or promptly treated. Sometimes these more serious symptoms develop without the individual ever having had a rash.

Reported Cases of Lyme Disease—United States, 2012

One dot is placed randomly within the county of residence for each confirmed case. Though Lyme disease cases have been reported in nearly every state, cases are reported based on the county of residence, not necessarily the county of infection.




Figure 1. Reported cases of Lyme disease in the United States, 2012. Though Lyme disease cases have been reported in nearly every state, cases are reported based on the county of residence, not necessarily the county of infection.

Credits: Centers for Disease Control



Figure 2. Black-legged deer tick female.

Credits: J. F. Butler, UF/IFAS

A different but similar disease occurs in Florida, and this disease is often confused with Lyme disease. Southern tick-associated rash illness is caused by a pathogen that is closely related to Lyme disease and vectored by the lone star tick. This disease was first reported in 2001 and occurs wherever lone star ticks are found. Similar to Lyme disease,

a red, expanding rash with a central, clear area at the site of the tick bite has been reported.

Treatment

Once diagnosed, Lyme disease can be treated. Physicians can determine if an individual has been infected by the Lyme disease organism using a simple blood test; however, some people test negative but have the disease. The CDC warns against unproven tests, and proper testing procedures are important.

Infection can be treated by taking certain antibiotics. However, no immunity is conferred from infection so a person could get Lyme disease again from another infected tick.

Pets and other animals can contract Lyme disease as well, exhibiting symptoms similar to those in humans. Veterinarians can test for Lyme disease in pets and domestic animals exhibiting suspicious signs of arthritis (in younger animals), heart problems, or neurological signs.

Antibiotics used for treatments of Lyme disease include doxycycline, amoxicillin, or cefuroxime axetil. When patients are treated early in the stages of infection they recover fast and completely. Patients diagnosed with later symptoms of the disease may have persistent or recurrent symptoms even when treated.

Prevention

The best prevention against Lyme disease is to avoid being bitten by ticks. Individuals who spend a lot of time outdoors should be aware of the danger and make it a habit to regularly check their bodies for ticks. The tick requires time to attach itself and begin feeding. It is possible to remove ticks before they are able to transmit the disease. Ticks should be grasped with tweezers at the point where their mouthparts enter the skin and pulled straight out with firm pressure. Immature ticks are small and difficult to detect; often they appear as a freckle or mole.

Stay out of dense undergrowth unless absolutely necessary. Walk on closely mowed grass or paved walkways whenever possible.

Wear long-sleeved shirts and long pants tucked into socks. Light-colored clothes make ticks more visible and easier to pick off.

Apply tick repellent to socks and shoes to prevent ticks from crawling onto clothing.

Inspect yourself and others thoroughly after walking through areas suspected of being infested with ticks.

When taking children on nature outings, keep them in a group with a leader who knows to avoid tick-infested areas and can inspect them for ticks.

See a doctor if Lyme disease symptoms appear.

A Lyme disease vaccine was withdrawn from the market, reportedly due to poor sales.

Tick Control

Recommendations for control of ticks include:

- Keep grass cut low to reduce humidity and thus kill ticks.
- Discourage wild animals (raccoons, skunks, deer, mice, etc.) from coming around your yard. They often harbor the ticks that transmit Lyme disease. Area treatment with insecticides is not warranted in most cases. However, in some situations it might be appropriate to use insecticides for control of ticks. Check with your local UF/IFAS Extension office for current UF/IFAS recommendations.

References

Anderson, JF, LA Magnarelli, W Burgdorfer, and AG Barbour. 1983. Spirochetes in *Ixodes dammini* and mammals in Connecticut. *Am. J. Trop. Med. Hyg.* 32: 818–24.

Barbour, AG and D Fish. 1993. The biological and social phenomenon of Lyme disease. *Science*. 260. 1610–16.

Burgdorfer, W, AG Barbour, SF Hayes, JL Benach, E Grunwaldt, and JP Davis. 1982. Lyme disease—a tick-borne spirochetosis? *Science*. 216: 1317–19.

Daniels, TJ, D Fish, JF Levine, MA Greco, AT Eaton, PJ Padgett, and DA LaPointe. 1993. Canine exposure to *Borrelia burgdorferi* and prevalence of *Ixodes dammini* (Acari: Ixodidae) on deer as a measure of Lyme disease risk in the northeastern United States. *J. Med. Entomol.* 30: 171–78.

Coyle, BS, GT Strickland, YY Liang, C Pena, R McCarter, and E Israel. 1996. The public health impact of Lyme disease in Maryland. *J. Infect. Dis.* 173: 1260–62.

Hayes, EB and J Piesman. 2003. How can we prevent Lyme disease? *N. Engl. J. Med.* 348: 2424–30.

Mather, TN, JMC Ribeiro, SI Moore, and A Spielman. 1988. Reducing transmission of Lyme disease spirochetes in a suburban setting. *Ann. New York Acad. Sci.* 539: 402–03.

Meek, JI, CL Roberts, EV Smith Jr, and ML Cartter. 1996. Underreporting of Lyme disease by Connecticut physicians. 1992. *J. Public. Health Manage. Pract.* 2: 61–65.

Pound, JM, JA Miller, JE George, and CA Lemeilleur. 2000. The '4-poster' passive topical treatment device to apply acaricide for controlling ticks (Acari: Ixodidae) feeding on white-tailed deer. *J. Med. Entomol.* 37: 588–94.

Schulze, TL, WM McDevitt, WE Parkin, and JK Shisler. 1987. Effectiveness of two insecticides in controlling *Ixodes dammini* (Acari: Ixodidae) following an outbreak of Lyme disease in New Jersey. *J. Med. Entomol.* 24: 420–24.

Schulze, TL, RA Jordan, and RW Hung. 1995. Suppression of subadult *Ixodes scapularis* (Acari: Ixodidae) following removal of leaf litter. *J. Med. Entomol.* 32: 730–33.

Schulze, TL, RA Jordan, RW Hung, RC Taylor, D Markowski, and MS Chomsky. 2001. Efficacy of granular deltamethrin against *Ixodes scapularis* and *Amblyomma americanum* (Acari: Ixodidae) nymphs. *J. Med. Entomol.* 38: 344–46.

Stafford, KC. 1991. Effectiveness of carbaryl applications for the control of *Ixodes dammini* (Acari: Ixodidae) nymphs in an endemic residential area. *J. Med. Entomol.* 28: 32–36.

Steere, AC, E Taylor, GL McHugh, and EL Logigian. 1993. The overdiagnosis of Lyme disease. *JAMA*. 269: 1812–16.

Wormser, GP, RB Nadelman, RJ Dattwyler, DT Dennis, ED Shapiro, AC Steere, TJ Rush, DW Rahn, PK Coyle, DH Persing, D Fish, and BJ Luft. 2000. Practice guidelines for the treatment of Lyme disease. *Clinic Infect. Dis.* 31: 1–14.