

Colletotrichum Crown Rot (Anthracnose Crown Rot) of Strawberry¹

Natalia A. Peres, M. S. Oliveira, and Steven J. MacKenzie²

Colletotrichum crown rot is caused by fungi in the *Colletotrichum gloeosporioides* species complex. These pathogens kill strawberry plants by aggressively invading crown tissue. Crown rot is a serious disease in warm production regions, such as those in the southeastern United States, since *C. gloeosporioides* grow and reproduce best at temperatures greater than 25°C (77°F). Although crown rot is observed in fields during the winter production season in Florida, it is most severe in nurseries in the southeastern United States and is one of the primary reasons that transplant production moved to cooler regions. Another species of *Colletotrichum*, *C. acutatum*, also causes plants to decline in vigor and wilt. Although it is sometimes isolated from crown tissue, the decline and wilt symptoms caused by this species are generally due to bud and root infection.

Causal Agent and Symptoms

The symptoms caused by three species within the *C. gloeosporioides* species complex that have been reported to affect strawberry in Florida (*C. siamense*, *C. theobromicola*, and *C. fructicola*) are virtually indistinguishable in the field. Plants infected initially show signs of water stress and subsequently collapse (Figure 1). This process may occur relatively rapidly, taking only 2 or 3 days at optimum temperatures (around 80°F). At lower temperatures (~68°F), plants may show initial signs of stress, but it could take several weeks before they collapse. Cutting through the crown tissue of infected plants lengthwise reveals a reddish-brown, firm rot (Figure 2). Infected plants from nurseries may grow normally (asymptomatic) for some time before symptoms occur. There are typically no lesions on foliage or stolons in production fields, although under greenhouse conditions or in summer nurseries, necrosis on stolons, lesions on fruit, or black leaf spots may be visible. Symptoms of Colletotrichum crown rot may be confused with those of *Phytophthora* crown rot (caused by *Phytophthora cactorum*) and charcoal rot (caused by *Macrophomina phaseolina*). To confirm a diagnosis, the pathogen must be isolated from the diseased crowns and identified in the laboratory.



Figure 1. Initial plant wilting symptoms of Colletotrichum crown rot.

Credit: UF/IFAS GCREC



Figure 2. Internal crown symptoms of Colletotrichum crown rot.

Credit: UF/IFAS GCREC

Disease Development and Spread

Plant propagation in Canada and northern states has greatly reduced the incidence of *Colletotrichum* crown rot. However, plants can become infected after transplant since *C. gloeosporioides* are abundant on noncultivated hosts in Florida, and genetic data indicate that some of this population are the same as those from strawberry crowns. Although inoculum from plants showing symptoms at the beginning of a production season can spread to other plants by the end of the season, the disease does not appear to be multicyclic and usually does not spread quickly from symptomatic to healthy plants. *Colletotrichum* species responsible for crown rot do not appear to survive between seasons on plant debris when plants are killed immediately after the production season ends in the spring, and the fungus disappears from crowns, plant residues, and soil during the hot and humid summer months in Florida.

There are no known cultivars that are immune to *Colletotrichum* crown rot, although cultivars do differ in susceptibility. The use of tolerant cultivars delays the onset of disease until later in the season or reduces the incidence of crown rot. Among the currently grown cultivars, 'Florida Beauty', and 'Florida Brilliance' are considered susceptible, whereas 'Sensation® Florida 127', 'Florida Ember' and 'Florida Encore' show moderate levels of resistance.

Disease Management

Using disease-free transplants and cultivar resistance are the most effective methods for controlling *Colletotrichum* crown rot in strawberry production fields. Currently, there is no certification program to guarantee that transplants are free of crown rot, and infected plants may not show symptoms until they have been established in the field. Transplants from northern latitudes or high-altitude nurseries in Canada and California should be disease-free since *C. gloeosporioides* is not present in these regions. Weekly foliar sprays of protectant fungicides such as captan are effective in suppressing the spread of crown rot from infected to healthy plants and reducing infections coming from native vegetation. However, protectant fungicides do not hinder the progress of symptoms in plants that are already infected. In this case, a single-site fungicide might be more effective, unless resistant strains are present in the field (see Table 1). Cultural practices that reduce the occurrence and movement of water on foliage, such as the use of drip irrigation, may limit the dispersal of the pathogen. Plants are also more sensitive to infection under high fertility conditions. Reduced nitrogen rates in nurseries or the use of nitrate rather than ammonium nitrogen sources may also reduce the risk of infection.

Table 1. Fungicides recommended for control of *Colletotrichum* crown rot of strawberries in Florida.

Product name (active ingredient)	Fungicide group ^a	Maximum rate/A		Minimum days to harvest	Re-entry interval (hours)	Remarks
		Per application	Per season			
Captan 80 WDG (captan)	M4	3.75 lb.	30 lb.	1	24	Do not apply in combination with or immediately before or closely following oil sprays. Do not mix with strongly alkaline materials.
Captec 4L® (captan)	M4	3 qt.	54 qt.	1	24	Do not apply in combination with or immediately before or closely following oil sprays. Do not mix with strongly alkaline materials.
Quilt Xcel® (propiconazole + azoxystrobin) ^b	3 + 11	14 fl. oz.	56 fl. oz.	0	12	Do not make more than two consecutive applications before alternating to another fungicide group and no more than four applications per crop per year.
Merivon (fluxapyroxad + pyraclostrobin) ^b	7 + 11	11 fl. oz.	33 fl. oz.	0	12	Do not make more than two sequential applications before alternating to another non-group 7 or non-group 11 fungicide
Abound® (azoxystrobin) ^b	11	15.4 fl. oz.	1.92 qt.	0	4	Do not make more than two consecutive applications and no more than four applications per crop per year.
Cabrio® EG (pyraclostrobin) ^b	11	14 fl. oz.	70 fl. oz.	0	12	Do not make more than two consecutive applications and no more than five applications per crop per year.
Evito® 480 SC, Aftershock™ (fluoxastrobin) ^b	11	5.7 fl. oz.	22.8 fl. oz.	1	12	Do not make more than two consecutive applications and no more than four applications per crop per year.
Flint® (trifloxystrobin) ^b	11	3.2 oz.	19.2 oz.	1	12	Do not make more than two consecutive applications and no more than six applications of group 11 fungicides per crop per year.

Switch® 62.5 WG (cyprodinil + fludioxonil)	9 + 12	14 oz.	56 oz.	0	12	Do not make more than two consecutive applications. Do not plant crops not on the label for 30 days after last application.
--	--------	--------	--------	---	----	--

^aFungicide group (FRAC Code): Numbers (1-37) and letters (M) are used to distinguish the fungicidal mode of action groups. All fungicides within the same group (with same number or letter) indicate same active ingredient or similar mode of action. This information must be considered when making decisions about how to manage fungicide resistance. M = multisite inhibitors, fungicide resistance is low; NC = not classified. Source: <https://www.frac.info/> (Fungicide Resistance Action Committee, FRAC).

^bResistance has been reported to group 11 fungicides in *C. gloeosporioides* isolates from Florida strawberry fields. Samples can be submitted to the UF/IFAS GCREC strawberry lab to determine if resistance is present.

¹ This document is PP 238, one of a series of the Department of Plant Pathology, UF/IFAS Extension. Original publication date August 2007. Revised September 2012, May 2013, December 2017, February 2022, and July 2025. Visit the EDIS website at <https://edis.ifas.ufl.edu> for the currently supported version of this publication.

² Natalia A. Peres, professor, strawberry pathology, Department of Plant Pathology, UF/IFAS Gulf Coast Research and Education Center; Michelle S. Oliveira, graduate student, Department of Plant Pathology, UF/IFAS Gulf Coast Research and Education Center; Steven J. MacKenzie, former research coordinator; Department of Plant Pathology, UF/IFAS Gulf Coast Research and Education Center; UF/IFAS Extension, Gainesville, FL 32611.

The Institute of Food and Agricultural Sciences (IFAS) is an Equal Opportunity Institution authorized to provide research, educational information and other services only to individuals and institutions that function with non-discrimination with respect to race, creed, color, religion, age, disability, sex, sexual orientation, marital status, national origin, political opinions or affiliations. For more information on obtaining other UF/IFAS Extension publications, contact your county's UF/IFAS Extension office. U.S. Department of Agriculture, UF/IFAS Extension Service, University of Florida, IFAS, Florida A & M University Cooperative Extension Program, and Boards of County Commissioners Cooperating. Andra Johnson, dean for UF/IFAS Extension.