

Bacterial Blight of *Ficus* Caused by *Xanthomonas*¹

Romina Gazis, Vanessa Campoverde and Aaron Palmateer²

Introduction

Florida leads the nation in the production of foliage plants, accounting for almost 70% of the national wholesale value. Species of *Ficus* are common ornamental foliage plants used for interior decoration and in home landscapes throughout warm climates in South Florida. *Ficus* species used as foliage plants include *F. altissima*, *F. benjamina*, *F. binneidijkii*, *F. elastica*, *F. lyrata*, *F. microcarpa*, *F. pumila*, *F. retusa*, and *F. rubiginosa*, among others (Chen, Henny, and McConnell 2002).

Symptoms

Xanthomonas species are bacterial plant pathogens that enter plants through wounds or natural openings, such as stomata (breathing pores) or hydathodes (pores through which the plant secretes water, often located at the leaf margin). Once the bacterium's cells enter the plant, they multiply and can move systemically, causing severe leaf blight. Initial symptoms appear as small, water-soaked, circular lesions with irregular borders (Figure 1). The water-soaked leaf areas can be readily observed by examining the underside of the leaves (Figure 2). After 7–14 days of initial symptoms, lesions enlarge, coalesce, and typically cover large portions of the leaf area (Figure 3). Eventually, the lesions turn brown with greenish-yellow borders, resulting in premature senescence and leaf drop (Figure 4–5).




Figure 1. *Xanthomonas* leaf spot on *Ficus elastica*. Left: Necrotic lesions are often distributed close to the main vein or the margin of the leaves.

Credits: Romina Gazis

Pathogen and Host Range

Xanthomonas campestris pathovar *fici* is reported to be the most common bacterial disease affecting *Ficus*. Host studies indicate that several *Ficus* species and their cultivars are susceptible to this pathogen, including *F. benjamina*, *F. buxifolia*, *F. triangularis*, *F. mexicana*, *F. maclellandii* 'Alli',

1. This document is PP305, one of a series of the Department of Plant Pathology, UF/IFAS Extension. Original publication date April 2013. Revised September 2024. Visit the EDIS website at <https://edis.ifas.ufl.edu> for the currently supported version of this publication.
2. Romina Gazis, associate professor, Department of Plant Pathology, and director, Plant Diagnostic Clinic, UF/IFAS Tropical Research and Education Center ; E. V. Campoverde, Extension agent II, UF/IFAS Extension Miami-Dade County; and A. J. Palmateer, courtesy professor of plant pathology, UF/IFAS TREC; UF/IFAS Extension, Gainesville, FL 32611.

The Institute of Food and Agricultural Sciences (IFAS) is an Equal Opportunity Institution authorized to provide research, educational information and other services only to individuals and institutions that function with non-discrimination with respect to race, creed, color, religion, age, disability, sex, sexual orientation, marital status, national origin, political opinions or affiliations. For more information on obtaining other UF/IFAS Extension publications, contact your county's UF/IFAS Extension office. U.S. Department of Agriculture, UF/IFAS Extension Service, University of Florida, IFAS, Florida A & M University Cooperative Extension Program, and Boards of County Commissioners Cooperating. Andra Johnson, dean for UF/IFAS Extension.

F. retusa ‘California Nitida’ and ‘Green Gem’, and *F. Grenn-island* (Chase and Henley 1993). The *Xanthomonas* isolates collected from diseased leaves of *F. elastica* in local nurseries (Homestead, Florida) have been tentatively identified as *Xanthomonas axonopodis*. Molecular characterization of the 16S rDNA exhibited 99% nucleotide identity to an isolate of *X. axonopodis* (GenBank accession No. AF123091).



Figure 2. Classic symptoms on older leaves showing on both sides of *Ficus* leaf.

Credits: Vanessa Campoverde




Figure 3. *Ficus elastica* plant inoculated with *Xanthomonas* showing disease progress. (A) Local lesions after 7 days. (B) Water-soaked lesions after 10 days. (C) Necrosis symptoms. (D) Death, 21 days after inoculation.

Credits: Vanessa Campoverde

Bacterial blight on *F. elastica* appears to affect plants at all stages and can kill young plants (i.e., liner stage). Host range studies conducted at the UF/IFAS Tropical Research

and Education Center (TREC) indicated that several economically important cultivars, including ‘Burgundy’, ‘Robusta’, and ‘Cabernet’, are equally susceptible to the disease.



Figure 4. *Xanthomonas* leaf spot on *Ficus elastica* from commercial nurseries. Advanced symptoms (coalescence of the necrotic lesions) are observed on the foliage.

Credits: Romina Gazis



Figure 5. High incidence and severity of *Xanthomonas* blight on *F. elastica*.

Credits: Ian Maguire, UF/IFAS

Disease Management

With bacterial diseases, a preventative disease management approach is the best option. The only way to confirm the presence of *Xanthomonas* blight on *Ficus* is to submit samples to a plant disease diagnostic laboratory. Contact your local UF/IFAS Extension agent for assistance. Successfully managing bacterial pathogens requires a long-term strategy and actions that focus on changing and improving

procedures and materials to reduce the opportunity for spread or reintroduction of the pathogen.

Monitor Incoming Plants

If buying from other locations, inspect new liners and plants upon delivery. Any new plants brought into the nursery should be kept isolated (also isolate tools and continuous bench space used for these new plants) from other plants for at least 3–4 weeks to observe any disease or pest symptoms and to avoid contamination with other crops. Do not forget about other potential contamination surfaces, like plant transport trailers or cart surfaces.

Scouting For Early Symptom Development

Bacteria are readily spread by wind, splashing water, contaminated bedding mix, diseased plants, and on the surface of materials (i.e., pots, pruning tools, etc.). Plant propagation is a particularly vulnerable type of production because young, tender plants and optimal propagation conditions favor disease development. If you suspect an unhealthy or diseased *Ficus*, contact your local UF/IFAS Extension agent for an accurate diagnosis.

Maintain a Disease-Free Environment

Sanitation is the key!

Disinfect tools when pruning plants. Bleach (sodium hypochlorite), which works by oxidizing or destroying the molecular bonds in fungi and lysing cells of bacteria, is used in a 10% concentration. Avoid mixing bleach with acids because toxic chlorine gas may result. Always use with good ventilation.

Store new pots in sanitized areas. Your best option is to always use new potting containers, but if this is not feasible, submerge potting containers in a 1:3 ratio of bleach to water with agitation for a minimum of 10 minutes.

Try to use only unopened potting soil and store on a covered, paved surface that can be periodically washed down with a 1:3 ratio of bleach to water.

Water Management

If possible, use a microjet or drip irrigation system. These systems do not allow water to touch the foliage, which can lead to bacterial spread.

Monitor irrigation and avoid watering late in the day or evenings. If using overhead irrigation, water in the morning so that the foliage has time to dry before evening. Allowing the foliage to dry reduces bacterial spread.

Chemical Control

Pesticides (i.e., fungicides and bactericides) are of limited value in managing bacterial diseases. However, they are your primary defense in prevention and, if used judiciously, they can be effective. Refer to the *Professional Disease Management Guide for Ornamental Plants* (<https://edis.ifas.ufl.edu/pp123>) for information about general pesticides used for managing diseases of ornamental plants.

Literature Cited

Campoverde, E. V., and A. J. Palmateer. 2011. "A Severe Outbreak of *Xanthomonas* on *Ficus elastica* in South Florida." *Proc. Fla. State Hort. Soc.* 124: 321–322.

Chase, A. R., and R. W. Henley. 1993. "Susceptibility of Some *Ficus* Species and Cultivars to *Xanthomonas*." *South-ern Nursery Digest* 27 (6): 20–21.

Chen, J., R. J. Henny, and D. B. McConnell. 2002. "Development of New Foliage Plant Cultivars." In *Trends in New Crops and New Uses*, edited by J. Janick and A. Whipkey, 466–472. Alexandria, VA: ASHS Press.