

# Cotinus coggygria 'Pendulus': 'Pendulus' Smoketree<sup>1</sup>

Edward F. Gilman, Dennis G. Watson, Ryan W. Klein and Deborah R. Hilbert<sup>2</sup>

### Introduction

This slow growing, open-crown, weeping, small tree eventually reaches to 10 to 15 feet tall (occasionally 20) and 10 to 15 feet wide. The large panicles of wispy cream-colored flowers produced in spring and early summer give the effect of a cloud of smoke. They make a wonderful accent in a shrub border and can be planted as a specimen or as a patio tree where the black, showy, multiple trunks can be displayed. Planting smoketree is a good way to extend the spring flowering-tree season into the summer before the crape-myrtles come into full bloom. Fall color is usually good to excellent and ranges from yellow to orange and brilliant red-purple. Many people grow it simply to enjoy the vivid fall color.

## **General Information**

Scientific name: Cotinus coggygria

**Pronunciation:** koe-TYE-nus koe-GUY-gree-uh

Common name(s): 'Pendulus' smoketree, 'Pendulus'

wig-tree, 'Pendulus' smokebush

Family: Anacardiaceae

**USDA hardiness zones:** 5B through 8B (Figure 2)

**Origin:** not native to North America




Figure 1. Middle-aged *Cotinus coggygria* 'Pendulus': 'Pendulus' smoketree.

**Invasive potential:** not assessed/incomplete assessment

**Uses:** specimen; deck or patio; container or planter; street without sidewalk; reclamation; trained as a standard; screen; parking lot island < 100 sq ft; parking lot island 100–200 sq ft; parking lot island > 200 sq ft; tree lawn 3–4 feet wide; tree lawn 4–6 feet wide; tree lawn > 6 ft wide; highway median

- 1. This document is ENH363, one of a series of the Department of Environmental Horticulture, UF/IFAS Extension. Original publication date November 1993. Revised December 2006 and December 2023. Visit the EDIS website at https://edis.ifas.ufl.edu for the currently supported version of this publication.
- 2. Edward F. Gilman, professor emeritus; Dennis G. Watson, former associate professor, Department of Agricultural and Biological Engineering; Ryan W. Klein, assistant professor, arboriculture; and Deborah R. Hilbert, UF/IFAS Gulf Coast Research and Education Center; Department of Environmental Horticulture; UF/IFAS Extension, Gainesville, FL 32611.

The Institute of Food and Agricultural Sciences (IFAS) is an Equal Opportunity Institution authorized to provide research, educational information and other services only to individuals and institutions that function with non-discrimination with respect to race, creed, color, religion, age, disability, sex, sexual orientation, marital status, national origin, political opinions or affiliations. For more information on obtaining other UF/IFAS Extension publications, contact your county's UF/IFAS Extension office.

U.S. Department of Agriculture, UF/IFAS Extension Service, University of Florida, IFAS, Florida A & M University Cooperative Extension Program, and Boards of County Commissioners Cooperating. Andra Johnson, dean for UF/IFAS Extension.




Figure 2. Range

# **Description**

Height: 10 to 15 feet

Spread: 10 to 18 feet

Crown uniformity: symmetrical

Crown shape: round, weeping

Crown density: dense

Growth rate: slow

Texture: medium

## **Foliage**

**Leaf arrangement:** alternate (Figure 3)




Figure 3. Foliage

**Leaf type:** simple

Leaf margin: entire

Leaf shape: obovate, elliptic (oval)

Leaf venation: reticulate, pinnate

Leaf type and persistence: deciduous

**Leaf blade length:** 2 to 4 inches

Leaf color: blue or blue-green, green

Fall color: yellow, orange, red, purple

Fall characteristic: showy

### **Flower**

**Flower color:** pink, white/cream/gray

Flower characteristics: very showy

### **Fruit**

Fruit shape: oval, irregular

Fruit length: less than 0.5 inch

Fruit covering: dry or hard

**Fruit color:** brown

**Fruit characteristics:** does not attract wildlife; showy; fruit/leaves not a litter problem

# **Trunk and Branches**

**Trunk/bark/branches:** branches droop; not showy; typically multi-trunked; thorns

Pruning requirement: needed for strong structure

Breakage: resistant

Current year twig color: brown

Current year twig thickness: medium, thick

Wood specific gravity: unknown

### **Culture**

Light requirement: full sun

Soil tolerances: clay; sand; loam; acidic; alkaline;

well-drained

**Drought tolerance:** high

Aerosol salt tolerance: moderate

### **Other**

Roots: not a problem

Winter interest: no

Outstanding tree: yes

Ozone sensitivity: unknown

Verticillium wilt susceptibility: susceptible

Pest resistance: resistant to pests/diseases

# **Use and Management**

The tree is tough and adapts to restricted soil spaces. It could be used along a street under power lines and would not require pruning for many years. It is a small tree, well-adapted to urban areas with almost year-round interest which should be used more in our landscapes.

Smoketree grows best in a sunny location and a well-drained loam. It will grow asymmetrically and lean toward the light in a partially sunny area, so it is best to locate it in full day sun where the crown will develop symmetrically. Though sometimes short-lived in rich soil, smoketree is useful in dry, rocky soil where there is no irrigation. It also grows in a wide range of soil pH, including alkaline. Probably short-lived (20 years—maybe more) in most situations but who cares—the tree is great while it's around!

Propagation of cultivars is by cuttings.

Cultivars include: 'Daydream'—heavy production of panicles, dense habit of growth; 'Flame'—brilliant orangered fall color, pink inflorescences; 'Nordine'—resembles 'Royal Purple'; 'Notcutt's Variety'—dark maroon-purple leaves (also known as 'Foliis Purpureis Notcutt's Variety' and 'Rubrifolius'); 'Purpureus'—leaves bronze-green, inflorescences purplish-pink; 'Royal Purple'—leaves are darker purple than 'Notcutt's Variety', rich red-purple fall color, purplish-red inflorescences; and 'Velvet Cloak'—dark purple-leaf form, spectacular fall color of reddish-purple.

#### **Pests**

Oblique-banded leaf roller mines and rolls the leaves, usually in June. The insect rarely occurs in sufficient numbers to warrant control. When it does occur, they can be picked off by hand or sprayed with *Bacillus thuringiensis*.

### **Diseases**

Leaf spots can be caused by various genera of fungi but are usually not serious.

Scab may cause leaf drop. No controls are listed.

The most serious disease is Verticillium wilt. Smoketree is very susceptible, and entire branches wilt and die. Prune out infected branches, fertilize, and water the plant during dry weather.

A stem canker can be a problem in the eastern US.