

Psidium cattleianum: Strawberry Guava¹

Edward F. Gilman, Dennis G. Watson, Ryan W. Klein, Andrew K. Koeser, Deborah R. Hilbert, and Drew C. McLean²

Introduction

This upright, multi-branched, evergreen shrub or small tree has many features which make it a popular landscape choice. Unfortunately, strawberry guava has escaped cultivation and has infested ditch banks and other wild areas in south Florida, and its use should be tempered. Capable of reaching 25 feet in height but often seen at 10 to 15 feet, the thick, smooth, dark green, four-inch-long, leathery leaves nicely complement the smooth, grey-brown to golden bark which attractively peels off in thin sheets. The single, white, one-inch-diameter flowers have many prominent stamens and look like powderpuffs in miniature. The blooms appear heaviest in April but can open sporadically throughout the year.

General Information

Scientific name: Psidium cattleianum

Pronunciation: SID-ee-um kat-lay-ee-AH-num **Common name(s):** strawberry guava, cattley guava

Family: Myrtaceae

USDA hardiness zones: 10A through 11 (Figure 2)

Origin: native to Brazil

UF/IFAS Invasive Assessment Status: invasive and not recommended (South); caution, may be recommended but manage to prevent escape (North, Central)



Figure 1. Full Form - *Psidium cattleianum*: strawberry guava Credits: UF/IFAS

Uses: shade; deck or patio; specimen; espalier; fruit; container or planter; highway median

- 1. This document is ENH-687, one of a series of the Environmental Horticulture Department, UF/IFAS Extension. Original publication date November 1993. Revised December 2018. Visit the EDIS website at https://edis.ifas.ufl.edu for the currently supported version of this publication.
- 2. Edward F. Gilman, professor emeritus, Environmental Horticulture Department; Dennis G. Watson, former associate professor, Department of Agricultural and Biological Engineering Department; Ryan W. Klein, graduate assistant, Environmental Horticulture Department; Andrew K. Koeser, assistant professor, Environmental Horticulture Department, UF/IFAS Gulf Coast Research and Education Center; Deborah R. Hilbert, graduate assistant, Environmental Horticulture Department, GCREC; and Drew C. McLean, biological scientist, Environmental Horticulture Department, GCREC; UF/IFAS Extension, Gainesville, FL 32611.

The Institute of Food and Agricultural Sciences (IFAS) is an Equal Opportunity Institution authorized to provide research, educational information and other services only to individuals and institutions that function with non-discrimination with respect to race, creed, color, religion, age, disability, sex, sexual orientation, marital status, national origin, political opinions or affiliations. For more information on obtaining other UF/IFAS Extension publications, contact your county's UF/IFAS Extension office. U.S. Department of Agriculture, UF/IFAS Extension Service, University of Florida, IFAS, Florida A & M University Cooperative Extension Program, and Boards of County Commissioners Cooperating. Nick T. Place, dean for UF/IFAS Extension.



Figure 2. Range

Description

Height: 15 to 25 feet **Spread:** 10 to 25 feet

Crown uniformity: irregular Crown shape: round, vase Crown density: moderate Growth rate: moderate Texture: medium

Foliage

Leaf arrangement: opposite/subopposite

Leaf type: simple **Leaf margin:** entire

Leaf shape: obovate, elliptic (oval) **Leaf venation:** pinnate, brachidodrome



Figure 3. Leaf - *Psidium cattleianum*: strawberry guava Credits: UF/IFAS

Leaf type and persistence: evergreen, broadleaf evergreen

Leaf blade length: 1 ½ to 4 inches

Leaf color: dark green on top, paler green underneath

Fall color: no color change **Fall characteristic:** not showy

Flower

Flower color: white

Flower characteristics: showy; fragrant; emerges singly or

in groups of 3 at leaf axis **Flowering:** year-round

Fruit

Fruit shape: round or obovoid Fruit length: 1 to 2 inches Fruit covering: fleshy berry Fruit color: purplish red

Fruit characteristics: attracts birds; showy; fruit/leaves a

litter problem

Fruiting: year-round



Figure 4. Fruit - *Psidium cattleianum*: strawberry guava Credits: UF/IFAS

Trunk and Branches

Trunk/branches: branches don't droop; showy; typically

multi-trunked; no thorns

Bark: gray to reddish brown, smooth, and peeling

Pruning requirement: little required

Breakage: resistant

Current year twig color: green, brown Current year twig thickness: thin, medium

Wood specific gravity: unknown



Figure 5. Bark - *Psidium cattleianum*: strawberry guava Credits: Gitta Hasing, UF/IFAS

Culture

Light requirement: full sun to partial shade **Soil tolerances:** clay; sand; loam; alkaline; acidic;

well-drained

Drought tolerance: moderate **Aerosol salt tolerance:** moderate

Other

Roots: not a problem Winter interest: no Outstanding tree: no

Ozone sensitivity: unknown

Verticillium wilt susceptibility: unknown **Pest resistance:** resistant to pests/diseases

Use and Management

The 1.5-inch-diameter, bright red, pear-shaped fruits ripen to a very dark red in July and are a popular treat - for people and wildlife. Birds, raccoons, and squirrels all love the delicious, sweet-tart, white flesh and will widely spread the small, grape-like seeds. The abundant, fallen fruits may be messy on hard surfaces such as walks and cars, and trees should probably not be planted along residential streets or over sidewalks or patios. It would make a good small boulevard or median street tree and accents an area with its well-shaped canopy and interesting trunk form. It combines well with other plants in a shrub border and provides both food and cover for wildlife.

Strawberry guava grows well in full sun or partial shade on a wide range of soils, including sand or clay, but fruit quality is improved on rich soils. Plants are drought-tolerant once established but benefit from a thick, organic mulch. Any necessary pruning should be done after fruiting. Pick the fruit for eating before the fruit fly larvae discover it.

Psidium littorale var. *littorale*, lemon guava, is slower-growing and has large yellow fruits with delightful, spicy flavor.

Propagation is by air-layering's or by seeds. Seeds germinate easily and rapidly. Seedling plants take seven or eight years to bear fruit.

Pests and Diseases

No pests or diseases are of major concern. Fruit fly larvae devour the fruit but do no harm to the tree.

Reference

Koeser, A.K., Friedman, M.H., Hasing, G., Finley, H., Schelb, J. 2017. Trees: South Florida and the Keys. University of Florida Institute of Food and Agricultural Sciences.