

# *Styphnolobium japonicum* 'Columnaris': Columnar Scholar Tree<sup>1</sup>

Edward F. Gilman, Dennis G. Watson, Ryan W. Klein, and Deborah R. Hilbert<sup>2</sup>

## Introduction

This cultivar of pagoda tree probably has a height of about 40 feet and spread of 15 to 20 feet, forming a fine-textured, upright canopy. It has a rapid growth rate and tolerates polluted city conditions, heat, and drought. The tree prefers a sunny, open location on any light soil. The very showy, greenish-white to yellow flowers are produced in mid- to late-summer and provide an airy feel to the tree for several weeks. A yellow dye can be made by boiling the dried flowers and buds in water. The young green twigs turn a dark grey with age. It has been reported that the species must be about 10-years-old to bloom, but the cultivar 'Regent' apparently blooms at an earlier age.

## General Information

**Scientific name:** *Styphnolobium japonicum*

**Pronunciation:** stiff-no-lobe-ee-um juh-PAWN-ih-kum

**Common name(s):** Columnar scholar tree, columnar Japanese pagoda tree

**Family:** *Fabaceae*

**USDA hardiness zones:** 5A through 8A (Figure 2)

**Origin:** not native to North America




Figure 1. Middle-aged *Styphnolobium japonicum* 'Columnaris': columnar scholar tree.  
Credits: UF/IFAS

**Invasive potential:** not assessed/incomplete assessment

**Uses:** specimen; street without sidewalk; sidewalk cutout (tree pit); tree lawn 3-4 feet wide; tree lawn 4-6 feet wide; tree lawn > 6 ft wide; urban tolerant; highway median

1. This document is ENH-751, one of a series of the Department of Environmental Horticulture, UF/IFAS Extension. Original publication date November 1993. Revised December 2006 and March 2024. Visit the EDIS website at <https://edis.ifas.ufl.edu> for the currently supported version of this publication.
2. Edward F. Gilman, professor emeritus; Dennis G. Watson, former associate professor, Department of Agricultural and Biological Engineering; Ryan W. Klein, assistant professor, arboriculture; and Deborah R. Hilbert, UF/IFAS Gulf Coast Research and Education Center; Department of Environmental Horticulture; UF/IFAS Extension, Gainesville, FL 32611.

The Institute of Food and Agricultural Sciences (IFAS) is an Equal Opportunity Institution authorized to provide research, educational information and other services only to individuals and institutions that function with non-discrimination with respect to race, creed, color, religion, age, disability, sex, sexual orientation, marital status, national origin, political opinions or affiliations. For more information on obtaining other UF/IFAS Extension publications, contact your county's UF/IFAS Extension office. U.S. Department of Agriculture, UF/IFAS Extension Service, University of Florida, IFAS, Florida A & M University Cooperative Extension Program, and Boards of County Commissioners Cooperating. Andra Johnson, dean for UF/IFAS Extension.




Figure 2. Range.  
Credits: UF/IFAS

## Description

**Height:** 35 to 45 feet

**Spread:** 15 to 20 feet

**Crown uniformity:** symmetrical

**Crown shape:** columnar, upright/erect

**Crown density:** moderate

**Growth rate:** moderate

**Texture:** fine

## Foliage

**Leaf arrangement:** alternate (Figure 3)




Figure 3. Foliage.  
Credits: UF/IFAS

**Leaf type:** odd-pinnately compound

**Leaf margin:** entire

**Leaf shape:** ovate

**Leaf venation:** pinnate

**Leaf type and persistence:** deciduous

**Leaf blade length:** less than 2 inches

**Leaf color:** green

**Fall color:** yellow

**Fall characteristic:** not showy

## Flower

**Flower color:** white/cream/gray, yellow

**Flower characteristics:** showy

## Fruit

**Fruit shape:** pod or pod-like, elongated

**Fruit length:** 3 to 6 inches, 6 to 12 inches

**Fruit covering:** dry or hard

**Fruit color:** brown, yellow

**Fruit characteristics:** does not attract wildlife; showy; fruit/leaves a litter problem

## Trunk and Branches

**Trunk/bark/branches:** branches don't droop; not showy; typically one trunk; thorns

**Pruning requirement:** needed for strong structure

**Breakage:** resistant

**Current year twig color:** green

**Current year twig thickness:** thin, medium

**Wood specific gravity:** unknown

## Culture

**Light requirement:** full sun

**Soil tolerances:** sand; loam; clay; acidic; alkaline; well-drained; occasionally wet

**Drought tolerance:** high

**Aerosol salt tolerance:** moderate

## Other

**Roots:** not a problem

**Winter interest:** no

**Outstanding tree:** no

**Ozone sensitivity:** unknown

**Verticillium wilt susceptibility:** unknown

**Pest resistance:** resistant to pests/diseases

## Use and Management

The tree drops flower petals creating a creamy white carpet for several weeks on the ground, but they can temporarily stain sidewalks. The yellow fruit pods form in late summer and are quite showy, dropping later in the winter and could be a nuisance to some people. But they are small and fairly easily washed away. The leaflets are small, creating light shade beneath the tree and are mostly washed away with rain or fall into shrub beds or between the grass blades.

Some trees come from the nursery with multiple trunks or branches clustered together at one spot on the trunk. Buy those with one central trunk growing up the center of the tree or prune the tree to a central leader to create a strong, durable structure. Space branches along the central leader to ensure good branch attachment. It may take several prunings to train the tree to the proper form.

This urban-tough tree is highly recommended for urban planting. It adapts to restricted soil spaces, tolerates salt spray, and tolerates drought in reasonable soil but not wet soil. Best when planted in full sun. This species has a few other cultivars: 'Pendula'—weeping habit; 'Princeton Upright'—upright form suitable for narrow sites, somewhat smaller than the species; 'Regent'—oval crown and blooms at an early age, has glossy leaves which shed soot and dirt, readily available in nurseries.

## Pests

Potato leafhopper kills young stems causing profuse branching or witches' broom on small branches. It usually is not a problem on larger trees.

## Diseases

*Styphnolobium* species is generally pest- and disease-free.

Occasionally, scholar tree will get a fungus canker about two-inches or less across, have raised reddish brown margins and light brown centers. The infected stem is killed when the fungus girdles the stem. Another fungus is sometimes found on dead branches on *Styphnolobium* species. Frost injury may give both fungi an entrance into the tree. Prune out dead, damaged, or diseased branches.

Twig blight or dieback can be a problem occasionally. Prune out infected branches and avoid unnecessary wounding. Keep trees vigorous by regular fertilization.

Powdery mildew forms a fungus mat which looks like a white coating on the leaves. The disease is usually not serious.