

# Florida's Bats: Velvety Free-Tailed Bat<sup>1</sup>

Holly K. Ober, Terry J. Doonan, and Emily H. Evans<sup>2</sup>

The velvety free-tailed bat (Figure 1) is found nowhere in the United States but extreme south Florida. With a wingspan of 11 inches (270 cm), it is a medium-sized bat. The color of its thick fur varies from dark brown to dark gray.



Figure 1. Velvety free-tailed bat (*Molossus molossus*).

Credits: Merlin Tuttle, Merlin Tuttle's Bat Conservation <http://www.merlintuttle.com/>

Found throughout the Greater and Lesser Antilles as well as much of Central and South America, the species has thus far been seen nowhere in the United States but the Florida Keys and Dry Tortugas (Figure 2). Their discovery in the United States was fairly recent: they were first found in the Florida Keys in the mid-1990s. There is some speculation they may occur in the Everglades region as well.




Figure 2. Velvety free-tailed bats in Florida are only found in the Florida Keys and Dry Tortugas.

Credits: Emily Evans

Velvety free-tailed bats in the United States are known to roost (sleep during the day) only in man-made structures. They have been found roosting in attics, crawl spaces, crevices, and under eaves in groups of as many as 1,300 individuals. In Cuba they also roost in tree hollows and in holes in utility poles.

1. This document is WEC380, one of a series of the Department of Wildlife Ecology and Conservation, UF/IFAS Extension. Original publication date October 2016. Reviewed October 2019. Visit the EDIS website at <https://edis.ifas.ufl.edu> for the currently supported version of this publication.
2. Holly K. Ober, associate professor and wildlife Extension specialist, Department of Wildlife Ecology and Conservation; Terry J. Doonan, mammal conservation coordinator, Florida Fish and Wildlife Conservation Commission; and Emily H. Evans, mammal conservation biologist, Florida Fish and Wildlife Conservation Commission; UF/IFAS Extension, Gainesville, FL 32611.

The Institute of Food and Agricultural Sciences (IFAS) is an Equal Opportunity Institution authorized to provide research, educational information and other services only to individuals and institutions that function with non-discrimination with respect to race, creed, color, religion, age, disability, sex, sexual orientation, marital status, national origin, political opinions or affiliations. For more information on obtaining other UF/IFAS Extension publications, contact your county's UF/IFAS Extension office. U.S. Department of Agriculture, UF/IFAS Extension Service, University of Florida, IFAS, Florida A & M University Cooperative Extension Program, and Boards of County Commissioners Cooperating. Nick T. Place, dean for UF/IFAS Extension.

These bats emerge from their roosts earlier than most other bats, often shortly before sunset. They feed on insects at high altitudes. It is believed that their large, forward-facing ears are aerodynamically aligned to conserve energy during flight.

## How do you know if you have found a velvety free-tailed bat?

First, check the rump to determine if a tail is visible. If you can see a tail more than an inch long (Figure 3), the bat is one of three species of free-tailed bats (in the family Molossidae).



Figure 3. Free-tailed bats (those in the family Molossidae) have long tails that extend more than an inch beyond their tail membrane, like the bat on the left. In comparison, other bats have short, stubby tails, like the bat on the right (not a free-tailed bat).

Credits: Merlin Tuttle, Merlin Tuttle's Bat Conservation <http://www.merlintuttle.com/>

Read below to determine how to differentiate the three species of free-tailed bats found in Florida.

### Velvety Free-Tailed Bat (*Molossus molossus*)

- Is found only in extreme south Florida (the Keys and perhaps the Everglades region)
- Typically roosts beneath roofs of man-made structures
- Roosts in the Keys in moderately-sized groups (50–300 individuals)
- Has ears that are not fully separated from one another (they are joined at the base; Figure 4)
- Each individual is about the size of a standard business card when it is at rest and its wings are folded

### Brazilian Free-Tailed Bat (*Tadarida brasiliensis*)

- Is found throughout the entire state of Florida, except the Keys
- Often roosts during the day in man-made structures (attics, stadiums, bridges, and bat houses)

- Typically roosts in large congregations (1,000–500,000 individuals)
- Produces a musky odor that is noticeable from some distance away
- Has ears that are fully separated from one another (not joined at the base like velvety free-tailed or bonneted bat ears; Figure 4)
- Is about the size of a standard business card when it is at rest and its wings are folded

### Florida Bonneted Bat (*Eumops floridanus*)

- Is found only in central and south Florida (as far north as Polk County)
- Roosts in pine tree cavities, palms, rock crevices, and man-made structures (beneath barrel tile roofs and in bat houses)
- Roosts in small groups (under 50 individuals)
- Has ears that are not fully separated from one another (they are joined at the base; Figure 4)
- Is the largest bat in Florida, about the size of a standard i-Phone when it is at rest and its wings are folded



Figure 4. The ears of the velvety free-tailed bat are joined at the base like the bat on the left, not separated like the Brazilian free-tailed bat on the right.

Credits: Kathleen Smith, FFWCC, used with permission; Elizabeth Braun de Torrez, UF/IFAS.

## Acknowledgement

This publication is funded in part by the Renewable Resources Extension Act (RREA).

## Additional Information

Florida Fish and Wildlife Conservation Commission.  
*Living with bats.*

Marks, C. S., and G. E. Marks. *Bats of Florida*. 2006. University Press of Florida. Gainesville, FL. 176 pp.

Ober, H. K., M. B. Main, and G. M. Allen. 2004. *Bats of Florida*. WEC186. Gainesville: University of Florida Institute of Food and Agricultural Sciences. <http://edis.ifas.ufl.edu/UW203>

Ober, H. K., and F. J. Mazzotti. 2008. *Conservation of Bats in Florida*. WEC247. Gainesville: University of Florida Institute of Food and Agricultural Sciences. <http://edis.ifas.ufl.edu/UW291>

Ober, H. K. 2008. *Effective Bat Houses for Florida*. WEC246. Gainesville: University of Florida Institute of Food and Agricultural Sciences. <http://edis.ifas.ufl.edu/uw290>

Ober, H. K. 2008. *Insect Pest Management Services Provided by Bats*. WEC245. Gainesville: University of Florida Institute of Food and Agricultural Sciences. <http://edis.ufl.edu/UW289>

Wisely, S. M., and H. K. Ober. 2015. *Facts about Wildlife Diseases: Rabies*. WEC239. Gainesville: University of Florida Institute of Food and Agricultural Sciences. <http://edis.ufl.edu/UW282>